

NGU Report 2014.057

Organic soil geochemistry in Nord-Trøndelag
and Fosen

| | | |
|---|---------------------------------|--|
| Report no.: 2014.057 | ISSN 0800-3416 | Grading: Open |
| Title: Organic soil geochemistry in Nord-Trøndelag and Fosen | | |
| Authors: Tor Erik Finne and Ola A. Eggen | | Client: NGU MINS |
| County: Nord-Trøndelag and Sør-Trøndelag | | Commune: |
| Map-sheet name (M=1:250.000) Trondheim, Østersund, Namsos, Grong, Vega | | Map-sheet no. and -name (M=1:50.000) |
| Deposit name and grid-reference: | | Number of pages: 82 Price (NOK): 315 NOK Map enclosures: |
| Fieldwork carried out: May-December 2013 | Date of report: Dec 19. 2014 | Project no.: 351700 Person responsible: <i>Brinda Tjern</i> |

Summary:

During field work in the summer and fall of 2013, organic soil samples were collected in a grid of 6 x 6 km in Nord-Trøndelag and Sør-Trøndelag's municipalities of Trondheim and Malvik as well as on the Fosen peninsula. Together with samples for quality control, the <2mm fraction of samples from 752 locations were digested by aqua regia and analyzed for 53 elements and Pb isotopes. The elements Be, In, Pd, Pt, Re and Te are kept out of the map collection due to very poor analytical data quality.

Results are documented with respect to quality of data and in tables of descriptive statistics, as well as plots of the cumulative probability function and by single element maps on a backdrop of bedrock geology.

| | | |
|------------------------|--------------|------------|
| Keywords: geochemistry | organic soil | aqua regia |
| MINS | | |
| | | |

CONTENTS

| | |
|---|----|
| 1. INTRODUCTION..... | 4 |
| 2. DESCRIPTION OF SURVEY AREA..... | 4 |
| 3. METHODS..... | 6 |
| 3.1 Planning stage and field work | 6 |
| 3.2 Sample preparation | 7 |
| 3.3 Chemical analyses | 8 |
| 3.4 Quality control | 9 |
| 3.4.1 Standard samples..... | 9 |
| 3.4.2 Duplicate samples | 10 |
| 3.4.3 Quality of Pb isotope analysis..... | 10 |
| 3.5 Data analysis..... | 11 |
| 4. RESULTS AND COMMENTS | 11 |
| 4.1 Data tables | 11 |
| 4.2 Cumulative probability (CP-) plot..... | 11 |
| 4.3 Maps | 12 |
| 4.4 First impressions..... | 14 |
| 5. ACKNOWLEDGEMENTS | 14 |
| 6. REFERENCES..... | 15 |
| Appendix 1: Cumulative frequency diagrams for all elements..... | 24 |
| Appendix 2: Maps of geochemistry in organic soil in Nord-Trøndelag and the Fosen peninsula..... | 33 |

FIGURES

| | |
|---|----|
| Figure 1: Area covered by previous and current studies..... | 5 |
| Figure 2:Map of quaternary deposits and ice movements in the surveyed area. | 6 |
| Figure 3:Photo showing sample station, five subsamples and some of the tools. Photo T.E. Finne..... | 7 |
| Figure 4: X-chart for P and Pb depicting stability for project standard Nmv. Dashed and dotted lines marks $\pm 10\%$ and $\pm 20\%$ deviation, respectively..... | 10 |
| Figure 5: The EDA symbol set. | 12 |
| Figure 6: Legend bedrock map | 13 |

TABLES

| | |
|---|----|
| Table 1: Overview of previous studies and methods | 8 |
| Table 2: Min., median, max. and precision values for the project humus standard Nmv..... | 17 |
| Table 3: Min., median, max. and precision values for the Acme standard CDV-1. | 18 |
| Table 4: Min., median, max. and precision values for the Acme standard VI6. | 19 |
| Table 5: Precision on analytical and field duplicates and also replicate analyses..... | 20 |
| Table 6: Statistical results (min., mean, max. and StDev) for the NIST standard | 21 |
| Table 7: Statistical results (min., 25%ile, median, mean, 75%ile, max., StDev and precision) for previous and present analyses of the Reimann et. al. (2007) study..... | 21 |
| Table 8: Statistical results (min., 25%ile, median, mean, 75%ile, max., StDev and precision) for previous and present analyses of the Jensen & Finne (2006) study | 21 |
| Table 9: Statistical parameters for the mapped data | 22 |
| Table 10: Comparison between Nord-Trøndelag & Fosen survey with Njåstad et. al. (1994), Ryghaug (1980) and Sæther (1985) | 23 |

1. INTRODUCTION

This survey of the organic soil (humus) covers the county of Nord-Trøndelag as well as adjacent parts of Sør-Trøndelag county on the Fosen peninsula and the municipalities of Trondheim and Malvik. The area was never mapped geochemically on a similar scale as was the case in the three northernmost counties of Norway in the 1980-ies, where stream sediments and stream water as well as mineral soil (Reimann et.al 2011) were common sampling media for all the counties. However, large parts of the area (below the tree line in Nord-Trøndelag county only) were sampled for organic soil characterization in 1960, reported by Ryghaug (1980) and by Finne and Grønlie (1983). Sæther (1985) collected several sampling media on a 1 sample/ca 30km² scale, among them organic soil. A national survey of humus chemistry containing 460-500 sampling stations throughout Norway was carried out in 1977, and repeated 1985 (Njåstad et.al, 1994) provides data on an even lower sampling density. The national survey turned into a monitoring program by being repeated 1995 and 2005 (Meyer et.al, 2104). A higher sampling density mapping of stream sediments (1 sample/3km²), covering Nord-Trøndelag and the Fosen peninsula, was carried out during the years 1983-1985 (Sæther, 1987, Sand, 1987, and Sæther, 1988). Figure 1 shows the extent of the different humus surveys.

This survey was planned and carried out in tandem with and in support of the mineral soil survey reported by Finne et.al (2014), and is a part of NGU's program MINS – Mineral resources in South Norway. In this report the term humus is used synonymous with organic soil, mostly for space considerations.

2. DESCRIPTION OF SURVEY AREA

The prominent topographical features of the Trondheimsfjord's major part and its arm Beitstadfjorden, as well as its continuation through the lake Snåsavatnet are all strong indicators of the geologic history of the area. The ENE-NE direction of these features coincides with major strike and with fracture zones like the Møre-Trøndelag Fault Zone (Gabrielsen and Ramberg, 1979; Nasuti et al, 2010). Along the coast are gneisses of the "Western Gneiss Region", 1850-1500 Myr, present to a varying extent, until they join similar rocks of the Transscandinavian Intrusive Belt crossing from the coast SE-wards through Lierne into Sweden (the "Grong culmination"). Most of the area, however, is covered by the rocks of the Caledonian nappes. According to the NGU ore database, ferrous metals dominate the precambrian rocks, whereas in the greenstones and schist of the Caledonian nappes, the base metals are dominant. The mining intensity, measured as number of records per area classified as "Mining" or Test mining" in the NGU ore database is for Nord-Trøndelag county 3.8/1000km², whereas the values for Nordland, Troms and Finnmark counties are 3.5, 3.4, and 1.0 respectively.




Figure 1: Area covered by previous and current studies.

The quaternary deposits of the area are dominated by areas of thin, discontinuous till material, interspersed with weathered rock of local origin. Figure 2 also show areas of till, mostly confined to lower altitudes in the mountain regions close to the Swedish border. Ice flow direction during the last glaciations is determined to be towards W-NW in general (Bargel et al, 1999), and is shown in the map as red points for each observation, with a line in the upstream direction.




Figure 2: Map of quaternary deposits and ice movements in the surveyed area.

3. METHODS

3.1 Planning stage and field work

Aiming for a sampling density similar to the surveys of the three northernmost counties, a grid of 6x6 km was generated to assist in planning of field logistics. Within each of the grid squares, field workers were free to find a suitable location, with a minimum distance of 10-100 m from abandoned to high traffic roads. The chosen site should, if possible, accommodate both sampling of the C-horizon mineral soil sample as well as the organic soil sample. Each sample of the organic soil layer was composed of a minimum of five subsamples taken from an area of about 50x50m, and from positions with a minimum of canopy cover. Subsamples were collected by cutting through vegetation and humus layer either with a steel cylinder of diameter 10cm or by a steel spade, forming a wedge or rectangle with the same area. A steel knife was then used to separate the 0-3 cm layer from the above vegetation and the deeper soil layers, ensuring a minimum of mineral soil material entering the Rilsan® plastic sample bag. Figure 3 shows a typical sample site above tree line, where the subsamples positioned throughout the sample locality, as well as some of the equipment. All locations were photographed, their positions recorded by handheld (WAAS/Egnos) GPS, and observations on humus thickness and vegetation characteristics were noted. Sample wet weight was on average 1.3 kg. Sample contamination was minimized by the field crew not wearing any jewelry during sampling, and tools were wiped clean before collecting the next sample. For

about every 20th sample a duplicate sample was collected 1-10m from original sample site, resulting in a total of 38 duplicate pairs. A total of 752 site/field samples were collected and accepted for further use, of these 95 were collected using helicopter for transport in remote areas.



Figure 3: Photo showing sample station, five subsamples and some of the tools. Photo T.E. Finne

3.2 Sample preparation

Upon arrival at the NGU laboratories, samples were dried in their original sampling bag for three weeks at temperatures below 40 °C. Sample dry weight was on average 0.55 kg. Subsequently all samples were dry sieved to <2mm (9 mesh), from which 2 aliquots of 1 dL were obtained and stored in Kautex polyethylene sample vial. Surplus <2mm material was stored in the original sample bag and saved for possible later usage. From all field duplicates, an additional split was generated.

Nylon sieves were used, and no jewelry was worn during preparation work. Cross contamination via sample dust during sieving was controlled by sieving samples one at a time in a vented box. All sieving equipment was cleaned using a vacuum cleaner in between every sample.

For the purpose of inserting a control sample in the analysis batch, two large volumes of organic soil were tested for homogeneity; a commercially available material from a garden

center and natural peat material removed from the residential lot of one of the authors. After drying, sieving to less than 2 mm and splitting, 20 1g samples of each of the two candidates were extracted by HNO₃ and analysed by ICP-AES. Following the same pattern, LOI was determined on 5g weighings heated to 480°C for 20 hours. The commercial soil was found to be the least homogenous, most likely due to its apparent added mineral soil mix of coarse (yet <2mm) material. The best choice material for control of stability of data was dubbed Nmv.

To facilitate use of this data set together with other data on acid extractable contents in humus, 15 samples from earlier investigations were included with this analytical batch. The preparation and analytical procedures differ somewhat between the data sets (see Table 1), but they nevertheless give an illustration of differences and similarities of these. The included samples were 10 of the 40 “LIT” samples of the 2005 GEOS transect (Reimann et.al, 2007), and 5 of the 42 samples collected in Finnmark 2005 (Jensen and Finne, 2006).

Following sample preparation, one series of all samples were randomized in a structured manner, so that for about every 17 samples sent to the laboratory, a field duplicate, its split and its ordinary sample as well as a split of the project standard Nmv were inserted. The control samples were not always inserted in the exact same positions within the group.

Table 1: Overview of previous studies and methods

| | | | |
|---|---|---|-----------------------------------|
| 39 control + 15 samples | Nmv (1 sample, 10 repetitions) | GEOS 2005 | Finnmark 2005 |
| N original, this survey | 10 & 39 Repetitions | 10 different samples, same 10 | 5 different, same 5 |
| Original: | | | |
| Sample weight, extraction, instrument | 1g HNO ₃ ICP-AES | 0,5 g HNO₃→Aqua Regia ICP-MS | 1g HNO ₃ ICP-AES |
| Reference material | None | 2x NIST1575a | 3x H-3 |
| This survey | | | |
| Method | 5g, HNO₃→Aqua Regia, ICP-MS | | |

3.3 Chemical analyses

The randomized series of 90+ g aliquots were shipped to ACME laboratories in Vancouver, Canada. The laboratory inserted further 28 splits of its own quality control (QC) sample CDV-1 for overall QC and 28 splits of the laboratory's own Pb isotope standard V16, as well as both of the certified reference materials NIST SRM 981 and NIST SRM 983 for Pb isotope QC. The laboratory also did replicate weighing, extraction and analyses of 28 replicate pairs throughout the analytical sequence.

A 5 g aliquot of the <2mm humus sample material was first leached with concentrated HNO₃ for 1 h and then digested in a hot (95 °C) water bath for an additional hour. After

cooling, an aqua regia solution of equal parts concentrated ACS grade HCl, HNO₃ and de-mineralised H₂O was added to each sample (6 mL/g solid) to leach in a hot (95 °C) water bath for 2 h. After cooling, the solution was made up to a final volume with 5% HCl and then filtered. The sample weight to solution volume ratio is 1 g per 20 mL. The solutions were analysed using a Spectro Ciros Vision emission spectrometer (ICP-AES) and a Perkin–Elmer Elan 6000 inductively coupled plasma mass spectrometer (ICP-MS) for a suite of 53 elements plus the Pb-isotopes 204, 206, 207 and 208.

Analytical results were returned from the laboratory within one month after receiving the samples. The remainder of the sample material was stored in the event of mishaps with the first weighing, and for possible upcoming analyses following alternative procedures. Unused sample material was not returned, but destroyed by the laboratory after the holding period, according to local regulations.

3.4 Quality control

To be able to estimate analytical precision based on analytical duplicates and to calculate the practical detection limits, it was agreed with the laboratory that all instrument readings were reported, independent of quantification and detection limits. For statistical calculations on the quality control part the instrument readings were used. Negative readings were replaced by a very low positive value prior certain statistical analyses.

Throughout the analysis there were inserted samples from certain previous studies for making it possible to do safe comparisons with these studies at a later stage. Results from these studies are not commented in this report.

3.4.1 Standard samples

X-charts are a simple but powerful way of studying the quality of the data. The data for a variable is plotted against its analytical sequence number, and by also plotting the median and deviation from the median it is possible to a) identify time trends or breaks in the analysis sequence, b) get an impression of precision by looking at the spread from the median, and c) get an impression of accuracy if the "true" or certified value is known.

X-charts from this survey indicate that no severe problems are present with regards to time trends or breaks in analytical results. Figure 4 serves an example where P and Pb in the MINS project organic soil standard Nmv are shown. The laboratory also used its own house standards, CDV-1 and V16, inserted throughout the analysis series. All in all, most results for the standards were satisfactory.




Figure 4: X-chart for P and Pb depicting stability for project standard Nmv. Dashed and dotted lines marks $\pm 10\%$ and $\pm 20\%$ deviation, respectively.

Values for minimum, median and maximum, as well as precision for the analytical results for the project standard Nmv and the laboratory standards CVD-1 and V16 are given in Table 2, Table 3 and Table 4, respectively.

3.4.2 Duplicate samples

Table 5 shows the estimate of precision based on the analytical duplicates and the field duplicates. The low precision is principally due to the natural variability shown in the difference between ordinary field sample and field duplicate samples. In most cases the observed problems with precision were due to very low concentrations as in the case of our project standard Nmv, i.e. analytical results at or below the limit of quantification, like Au, Be, Pd, Pt, Re and Te. In addition, the field duplicate results reveal that also elements In, V and W are plagued by poor reproducibility, and maps should be viewed with care.

Practical detection limit (PDL) was established based on method described by Demetriadis (2011), using the results for analytical replicates. Good results of the replicates led us to use a lower PDL rather than the laboratory's method detection (reporting) limit (MDL) for B, S, Sn, Th, Tl, V, W. On the other hand, PDL had to be increased for a range of elements; As, Au, Be, Cd, Ge, Hf, Li, Pd, Pt, Sc, Se, Te.

3.4.3 Quality of Pb isotope analysis

28 samples of both the NIST SRM 981 common lead isotopic standard (NIST.gov) and NIST SRM 983 radiogenic lead isotopic standard (NIST.gov) was inserted by the lab to the analysis series. The results (see Table 6) shows satisfactory accuracy for the SRM 981 and SRM analyses. The precision is not as satisfactory compared to the certified documentation, but we consider this to be less importance.

Among the inset of samples described in Table 1 Pb isotope data already existed for the Reimann et. al (2007) and Jensen & Finne (2006) studies. Despite differences in sample weight, extraction, and instrumentation the present study stands a comparison with these existing data. Table 7 and Table 8 shows a comparison of statistical parameters for the Reimann et. al (2007) and Jensen & Finne (2006) studies, respectively.

3.5 Data analysis

Geochemical data are compositional data, meaning that they do not contain truly independent values but only relative information; the reported concentrations of all elements analyzed depend on one another (Aitchison, 1986; Filzmoser et al., 2009). Such data have some special properties which can lead to wrong results when applying the methods developed for classical statistical data analysis (Reimann et al., 2013). Thus EDA (exploratory data analysis) techniques and simple order statistics as suggested by Reimann et al. (2008) are used here. All statistical calculations are determined by use of the freely available R software (R development core team, 2014) and the additional StatDA package (Filzmoser, 2013).

4. RESULTS AND COMMENTS

4.1 Data tables

A statistical overview for the dataset is provided in Table 9. The table is built around the minimum, maximum and median value, and also provides the values for a number of additional quantiles (percentiles) for the analyzed elements. As an additional measure of variation the “powers” are provided, which provide a direct impression of the orders of magnitude variation for each variable. When using classical statistical methods for calculation of the mean and standard deviation to derive at “thresholds” for anomalies, 2.6% of all data is often identified as anomalies at both ends of the distribution if the dataset has a normal distribution. The data at hand are far from normally distributed and therefore unsuited for classical statistics – thus the quantiles Q2 and Q98 (or Q5 and Q95) can be taken as lower and upper threshold for the data. However, quite often Cumulative Probability (CP) plots (see below) provide a better means of identifying anomalies in the data by inspection of shape of the curve.

Table 10 displays the analytical results with a more common approach, showing median, 98th percentile and maximum concentration for the organic soil dataset and data for the comparable 1985 national organic soil survey reported in Njåstad et.al (1994) dataset. For median, Q98 and maximum, the highest value between the four datasets is underlined. In the same table, Table 10 statistics for two other organic soil surveys by Sæther (1985) and by Ryghaug (1980) are also given. Sampling procedure for the latter called for 10 subsamples. These studies covers only parts of the MINS area, see Figure 1. All three datasets compared to the present are based on < 2mm humus ashed at 450-80 °C prior to HNO₃ extraction and ICP-AES analysis.

4.2 Cumulative probability (CP-) plot

Plots of the cumulative distribution function are one of the most informative displays of geochemical distributions (Reimann et al., 2008). In the plots the concentration is plotted along the X-axis and the cumulative probability is plotted along the Y-axis, and it allows the direct visual recognition of breaks in the curve which may be indicative of different geochemical processes. Breaks in the uppermost few percentiles of the distribution are often used as thresholds for anomaly identification. Readings below the PDL are here set to half the

PDL value for that element, respectively. Appendix 1 provides the CP-plots for all 53 variables and four Pb isotopes.

4.3 Maps

Many different methods for producing geochemical maps exist (see discussion in Reimann, 2005 or in Chapter 5 of Reimann et al., 2008). In mineral exploration so called “growing dot maps” as introduced by Bjørklund and Gustavsson (1987) are probably most often used. However, they focus the attention almost exclusively on the high values, the “anomalies” and are less well suited to study the data in more detail, e.g., in relation to geology. It may also be argued that the "growing dot map" has limitations in detecting local anomalies as they often do not display especially high values in relation to the whole dataset, but rather high values in relation to their local surroundings. Some of these shortcomings can be helped by giving special attention to the growth increment of the symbols, and the overall size of the symbols in the map image.

| | EDA symbol set | EDA symbol set extremes are accentuated | EDA symbol set extremes are accentuated Used in this report | Percentiles used in this report |
|----------------|----------------|--|---|------------------------------------|
| Highest values | + | ■ | ■ | 95–100% |
| Higher values | + | + | ● | 75–95% |
| Inner values | . | . | . | 25–75% |
| Lower values | ○ | ○ | ○ | 5–25% |
| Lowest values | ○ | ○ | ○ | 0–5% |

Figure 5: The EDA symbol set.

The EDA symbol set aim to provide an optical weight for each symbol in the map (Reimann et. al, 2008). Lower values are shown by circles, the inner (most common and in many cases the "least interesting") values are shown as dots, while the higher values are shown by crosses in the original EDA symbol set. Figure 5 shows the original EDA symbols to the left, and modifications in the middle. The percentiles used for the classes are 5 – 25 – 75 – 95%. All the maps are prepared on a backdrop of a generalized bedrock map based on the available maps in scale 1:250 000 hosted by <http://geo.ngu.no/kart/berggrunn/>. An excerpt of the legend for the 1:250 000 scale map series is shown in Figure 6. All geochemical maps are provided in Appendix 2. Please note that elements Be, In, Pd, Pt, Re and Te are kept out of the map collection due to very poor analytical data quality.

The dataset for this report is provided online

(<http://www.ngu.no/en-gb/tm/About-NGU/Projects/Mineralressurser-i-Sør-Norge-MINS>

Look for “Last ned data her”), and it is therefore possible and up to the reader to use different mapping techniques. Note, however, that in the provided data file all values below detection are marked as “<n”, n being the PDL, while NGU had the original instrument readings

available, i.e. values for every sample. NGU used the instrument reading values as these results often contain valuable information when using large datasets with hundreds of samples. For example, the laboratory's official detection limit for B is 1 mg/kg, but the QC results indicate that values down to 0.5 mg/kg are still reliable. Thus 10% real, natural variation would have been lost when setting all values below the MDL to for instance ½ of the detection limit.

| Berggrunn tegnforklaring | Bedrock legend |
|--|---|
| 2 - Sandstein | 2 - Sandstone |
| 3 - Konglomerat, sedimentær breksje | 3 - Conglomerate, sedimentary breccia |
| 7 - Sedimentære bergarter (uspesifisert) | 7 - Sedimentary rock (unspecified) |
| 8 - Skifer, sandstein, kalkstein | 8 - Shale, sandstone, limestone |
| 9 - Sandstein, skifer | 9 - Sandstone, shale |
| 10 - Kalkstein, skifer, mergelstein | 10 - Limestone, shale, marlstone |
| 11 - Kalkstein, dolomitt | 11 - Limestone, dolomite |
| 21 - Granitt, granodioritt | 21 - Granite, granodiorite |
| 22 - Dioritt, monzodioritt | 22 - Diorite, monzodiorite |
| 23 - Syenitt, kvartssyenitt | 23 - Syenite, quartz syenite |
| 24 - Monzonitt, kvartsmonzonitt | 24 - Monzonite, quartz monzonite |
| 25 - Mangerittsyenitt | 25 - Mangerite syenite |
| 26 - Rylolitt, ryodacitt, dacitt | 26 - Rhyolite, rhyodacite, dacite, keratophyre |
| 29 - Vulkanske bergarter (uspesifisert) | 29 - Volcanic rocks (unspecified) |
| 30 - Mangeritt til gabbro, gneis og amfibolitt | 30 - Mangerite to gabbro, gneiss and amphibolite |
| 35 - Gabbro, amfibolitt | 35 - Gabbro, amphibolite |
| 38 - Kvartsdioritt, tonalitt, trondhjemitt | 38 - Quartz diorite, tonalite, trondhjemite |
| 40 - Olivinstein | 40 - Peridotite, pyroxenite |
| 41 - Eklogitt | 41 - Eclogite |
| 45 - Anortositt | 45 - Anorthosite |
| 50 - Amfibolitt og glimmerskifer | 50 - Amphibolite and mica schist |
| 55 - Grønnstein, amfibolitt | 55 - Greenstone, amphibolite |
| 60 - Metasandstein, skifer | 60 - Metasandstone, mica schist |
| 61 - Kvartsitt | 61 - Quartzite |
| 62 - Glimmergneis, glimmerskifer, metasandstein, amfibolitt | 62 - Mica gneiss, mica schist, metasandstone, amphibolite |
| 65 - Fyllitt, glimmerskifer | 65 - Phyllite, mica schist |
| 66 - Kalkglimmerskifer, kalksilikatgneis | 66 - Calcareous mica schist, calc silicate gneiss |
| 70 - Marmor | 70 - Calcite marble |
| 71 - Dolomitt | 71 - Dolomite marble |
| 82 - Diorittisk til granittisk gneis, migmatitt | 82 - Dioritic to granitic gneiss, migmatite |
| 85 - Øyegneis, granitt, foliert granitt | 85 - Augen gneiss, granite, foliated granite |
| 87 - Båndgneis (amfibolitt, hornblendegneis, glimmergneis), stedvis migmatittisk | 87 - Amphibolite, hornblende gneiss, mica gneiss locally migmatitic |

Figure 6: Legend bedrock map

4.4 First impressions

As the main purpose of this report is to make the data publicly available together with a quality description, only some brief comments on the results will be given. When results of ongoing analysis of loss on ignition, pH, total carbon and total sulphur and nitrogen are reported, the data will be interpreted to a greater depth.

A few of the element maps outline distinct geographic features that to some degree is easily explained by other processes than those of interest for the ore prospector. Sodium, Na, is the most obvious, forming a continuous high value region along the coast. The marine influence through precipitation is seen for about 40 km distance from the coast. Mercury, Hg, on the other hand, does not show nearly the same continuous high level in the same area, and has numerous single point high values far from the coast, as well as some clusters of values above the 75%ile associated with a > 95%ile value. These are mostly located within the Caledonian schists or greenstones. Tin (Sn) is another element that demonstrates larger areas of high values. By first glance, one could think that also Sn was of marine origin, it should, however, rather be focused on the fact that most of the values are associated with granites or granitic gneisses.

Grongfeltet, known for its numerous base metal sulfide mineralizations, has several locations of high manganese concentrations, some of which are accompanied by high copper (and zinc) values. The samples in this area with high Cu-Zn concentrations may indicate real anomalies, and could easily be checked against data from the parallel mineral soil survey and similar data from the well explored area.

5. ACKNOWLEDGEMENTS

Fylkesmannen Nord-Trøndelag and the boards of all national parks in the area kindly accepted our application for “scientific investigation” in all the national parks – important when the whole picture is needed to understand its details. The municipalities that were asked, and the Blåfjella-Skjærkjellene national park board all granted legal provisions for use of helicopter, and a number of land owners kindly let us land on their properties. Guri Kjesbu at Værdalsbruket AS kindly gave car access to Juldalen, and we are also indebted to the landowner giving access to a military escorted road to Geitfjellet, Grong. We greatly appreciate the cooperative spirit of the local authorities and population; without it we could have risked the onset of snow before the field work was completed. The field crew did a formidable job: NGU's Malin Andersson, Belinda Flem, Guri Venvik Ganerød, Henning K. B. Jensen, Øystein Jæger, Agnes M. Raaness, Clemens Reimann, Anna Seither and Ola Vikhammar, as well as the authors. Jostein Jæger and Iselin Esp Pettersen worked relentlessly at the sample preparation lab by sieving, splitting and weighing the samples.

6. REFERENCES

- Aitchinson, 1986. The statistical analysis of compositional data. Chapman and Hall, London.
- Bargel, T., Huhta, P., Johansson, P., Lagerbäck, R., Mäkinen, K., Nenonen, K., Olsen, L., Rokoengen, K., Svedlund, J.-O., Väänänen, T. & Wahlroos, J.-E., 1999: *Maps of Quaternary geology in Central Fennoscandia, sheet 3: Ice-flow Indicators, scale 1:1 000 000, and Quaternary stratigraphy, scale 1:2 000 000*. Geological Surveys of Finland (Espoo), Norway (Trondheim) and Sweden (Uppsala).
- Bjørklund and Gustavsson, 1987. *Visualization of geochemical data on maps: new options*. J. of Geochem. Exploration, vol 29, pp. 89-103
- Demetriades, A., 2011. *Understanding the Quality of Chemical Data from the urban environment - Part 2: Measurement Uncertainty in the Decision-making Process*. In *Mapping the Chemical Environment of Urban Areas*. Johnson, C.C., Demetriades, A., Locutura, J. and Ottesen, R.T. (Eds.). Wiley-Blackwell, Oxford, united Kingdom. ISBN 978-0-470-74724-7
- Filzmoser P., Hron, K., Reimann, C., 2009. *Univariate statistical analysis of environmental (compositional) data – Problems and possibilities*. Science of the Total Environment 407, 6100-6108.
- Filzmoser, 2013. StatDA: Statistical Analysis for Environmental Data. R package version 1.6.7, <http://CRAN.R-project.org/package/StatDAR> core
- Finne, T.E. and Grønlie, A. 1983. *Humusgeokjemi i Nord-Trøndelag. En videre bearbeiding av data fra Landsskogtakseringen 1960*. NGU rapport 1889/2 (in Norwegian). 9 pp.
- Finne, T.E., Reimann, C., and Eggen, O.A. 2014. *Mineral soil geochemistry in Nord-Trøndelag and Fosen*. NGU report 2014.047. 91 pp.
- Jensen, H.K.B. and Finne, T.E. 2006. *Temporal variations 1995 - 2005 for heavy metals, arsenic and PAH in the humus layer in Pasvik and Jarfjord, Finnmark County, Norway*. NGU report 2006.017. 22 pp.
- Gabrielsen, R.H. & Ramberg, I.H. 1979: *Fracture pattern in Norway from Landsat imagery: Results and potential use*, Proceeding of the Norwegian Sea Symposium, Tromsø, Norwegian Petroleum Society. NSS 23, 1-28
- Meyer, M., Pesch, R., Schröder, W., Steinnes, E., and Uggerud, H.T. 2014. *Spatial patterns and temporal trends of heavy metal concentrations in moss and surface soil specimens collected in Norway between 1990 and 2010*. Environmental Sciences Europe 2014, 26:27
- Nasuti, A., Beiki, M & Ebbing, J. 2010: *Gravity and data acquisition over a segment of the Møre-Trøndelag Fault Complex*, NGU report 2010.049, 1-48
- NIST SRM 981 Certificate of Analysis, <https://www-s.nist.gov/srmors/certificates/981.pdf>
Retrieved Sep. 23 2014
- NIST SRM 983 Certificate of Analysis, <https://www-s.nist.gov/srmors/certificates/983.pdf>

Retrieved Sep. 23 2014

Njåstad, O., Steinnes, E., Bølviken, B., Ødegård, M. 1994. *Landsomfattende kartlegging av elementsammensetning i naturlig jord: Resultater fra prøver innsamlet i 1977 og 1985 oppnådd ved ICP emisjonsspektrometri*. NGU rapport 94.027 (in Norwegian). 114 pp.

Reimann, C., Arnoldussen, A., Englmaier, P., Filzmoser, P., Finne, T.E., Garrett, R.G., Koller, F. and Nordgulen, Ø. 2007. *Element concentrations and variations along a 120-km transect in southern Norway – Anthropogenic vs. geogenic vs. biogenic element sources and cycles*. Applied Geochemistry Vol. 22-4, pp 851-871.

Reimann, C., Filzmoser, P., Garrett, R.G., Dutter, R., 2008. Statistical data analysis explained. Applied environmental statistics with R. Wiley, Chichester, U.K.

Reimann, C., Birke, M., Demetriades, A., Filzmoser, P. & O'Connor, P. (eds.) (2013): Chemistry of Europe's Agricultural Soils. Part A: Methodology and interpretation of the GEMAS Data Set.-Geol. Jb., B 102,: 528 pp: 358 figs., 86 Tables, 1 DVD; Hannover

Ryghaug, P. 1980. *Geokjemisk undersøkelse av skogjorda i Nord-Trøndelag i forbindelse med Landsskogtakseringens markarbeid sommeren 1960*. NGU rapport 402 (in Norwegian). 22 pp.

Sand, K. 1987. *Bekkesedimenter fra Nord-Trøndelag og Fosenhalvøya. En geokjemisk undersøkelse av en ikke-magnetisk tungmineralfraksjon*. NGU rapport 87.057 (in Norwegian). 77 pp.

Sæther, O.M. 1985. *Geokjemi i Nord-Trøndelag - humus i områdene vest for riksvei E6*. NGU rapport 85.168 (in Norwegian). 46 pp.

Sæther, O.M. 1987. *Geokjemi i Nord-Trøndelag - en regional oversikt. Rapport med 1 vedlegg*. NGU rapport 87.082 (in Norwegian). 120 pp.

Sæther, O.M. 1988. *Geokjemiske undersøkelser i Nord-Trøndelag nøytronaktiviseringsanalyse på bekkesedimenter*. NGU rapport 88.108 (in Norwegian). 86 pp.

Table 2: Minimum, median, maximum and precision values for the MINS project organic soil standard Nmv. Concentrations in mg/kg.

| NmV standard (n=38) alphabetical | | | | | | | | | Sorted by precision | | |
|----------------------------------|-----------|-------|-------|---------|-------------------|---------|--------|---------|---------------------|-----------|---------|
| Element | Precision | | | Element | Precision | | | Element | Precision | | Element |
| | Min | Q50 | Max | | Min | Q50 | Max | | Precision | Precision | |
| Ag | 0,0 | 0,0 | 0,1 | 13,6 | Ni | 11,4 | 12,7 | 15,6 | 9,0 | Pt | 1286 |
| Al | 5767 | 6320 | 7348 | 7,1 | P | 383 | 424 | 463 | 4,8 | In | 309 |
| As | 2,23 | 2,91 | 3,66 | 11,70 | Pb | 4,84 | 5,36 | 6,13 | 4,92 | Pd | 219 |
| Au | -0,001 | 0,001 | 0,014 | 67,5 | ²⁰⁴ Pb | 0,1 | 0,1 | 0,1 | 5,1 | Te | 94 |
| B | 1,8 | 2,4 | 3,8 | 18,0 | ²⁰⁶ Pb | 1,2 | 1,3 | 1,5 | 5,9 | Au | 68 |
| Ba | 36 | 44 | 78 | 13,90 | ²⁰⁷ Pb | 1,02 | 1,15 | 1,29 | 5,46 | Re | 63 |
| Be | 0,0 | 0,1 | 0,3 | 52,0 | ²⁰⁸ Pb | 2,5 | 2,8 | 3,2 | 5,2 | Be | 52 |
| Bi | 0,0 | 0,0 | 0,1 | 11,7 | Pd | -0,0056 | 0,0014 | 0,0087 | 219,0 | Ge | 46 |
| Ca | 17607 | 19966 | 22241 | 3,48 | Pt | -0,0023 | 0,0001 | 0,0025 | 1286,00 | Ta | 33 |
| Cd | 0,1 | 0,1 | 0,2 | 15,5 | Rb | 4,4 | 5,0 | 5,7 | 5,7 | Sn | 25 |
| Ce | 6,5 | 7,0 | 7,8 | 3,1 | Re | 0,0000 | 0,0034 | 0,0092 | 63,1 | W | 24 |
| Co | 4,65 | 5,85 | 6,87 | 12,70 | S | 1655 | 3592 | 4369 | 9,99 | Hg | 18 |
| Cr | 19,1 | 22,2 | 28,6 | 10,0 | Sb | 0,1 | 0,1 | 0,2 | 10,9 | B | 18 |
| Cs | 0,5 | 0,5 | 0,6 | 4,1 | Sc | 1,4 | 1,7 | 2,0 | 12,5 | Hf | 18 |
| Cu | 19,7 | 23,3 | 31,6 | 8,14 | Se | 0,54 | 0,94 | 1,34 | 12,70 | Tl | 16 |
| Fe | 8316 | 9413 | 11161 | 5,9 | Sn | 1,2 | 2,0 | 17,3 | 25,3 | Cd | 16 |
| Ga | 1,3 | 1,6 | 1,8 | 8,7 | Sr | 34,3 | 37,2 | 45,2 | 5,6 | Ba | 14 |
| Ge | -0,02 | 0,06 | 0,12 | 46,30 | Ta | 0,004 | 0,008 | 0,014 | 33,40 | Ag | 14 |
| Hf | 0,0 | 0,1 | 0,1 | 17,6 | Te | -0,02 | 0,02 | 0,13 | 94,3 | Co | 13 |
| Hg | 0,0 | 0,0 | 0,1 | 18,4 | Th | 0,45 | 0,55 | 0,70 | 8,6 | Se | 13 |
| In | -0,03 | 0,00 | 0,02 | 309,00 | Ti | 441 | 524 | 676 | 9,45 | Sc | 13 |
| K | 790 | 863 | 959 | 5,2 | Tl | 0,029 | 0,038 | 0,050 | 16,3 | As | 12 |
| La | 2,9 | 3,3 | 3,8 | 6,3 | U | 0,22 | 0,26 | 0,29 | 5,4 | Bi | 12 |
| Li | 2,89 | 3,61 | 4,77 | 9,90 | V | 5,63 | 12,30 | 17,70 | 11,70 | V | 12 |
| Mg | 3554 | 3921 | 5002 | 8,1 | W | 0,03 | 0,08 | 0,11 | 23,7 | Sb | 11 |
| Mn | 174 | 192 | 221 | 4,7 | Y | 1,9 | 2,1 | 2,3 | 4,1 | Cr | 10 |
| Mo | 0,57 | 0,66 | 0,76 | 7,27 | Zn | 29,8 | 36,2 | 43,2 | 6,88 | S | 10 |
| Na | 129 | 146 | 183 | 8,0 | Zr | 2,6 | 2,9 | 3,5 | 4,9 | Li | 9,9 |
| Nb | 0,5 | 0,5 | 0,6 | 8,2 | | | | | | Ti | 9,5 |

Table 3: Minimum, median (Q50), maximum and precision values for the Acme standard CDV-1. Concentrations in mg/kg

| Acme standard CDV-1 (N=28), alphabetical | | | | | | | Sorted by precision | | | | | | |
|--|------------|---------|---------|-----------|---------|------------|---------------------|---------|-----------|---------|-----------|---------|-----------|
| Element | Min | Q50 | Max | Precision | Element | Min | Q50 | Max | Precision | Element | Precision | Element | Precision |
| Ag | 0,00783 | 0,0109 | 0,025 | 29 | Ni | 6,44 | 7,15 | 7,78 | 5,2 | Pt | -668 | Hg | 8,2 |
| Al | 1536 | 1706 | 1914 | 5,1 | P | 387 | 415 | 457 | 5,4 | Pd | -317 | Pb206 | 8,2 |
| As | 0,861 | 1,52 | 1,83 | 17 | Pb | 1,05 | 1,15 | 1,33 | 8,2 | Re | 148 | Zr | 7,6 |
| Au | 0,00138 | 0,00266 | 0,005 | 25 | Pb204 | 0,0119 | 0,0152 | 0,017 | 8,6 | Ge | 126 | Na | 7,2 |
| B | 9,84 | 12,6 | 14,3 | 9,0 | Pb206 | 0,225 | 0,291 | 0,332 | 8,2 | In | 99 | Ga | 7,1 |
| Ba | 9,18 | 9,86 | 11,4 | 3,9 | Pb207 | 0,206 | 0,239 | 0,268 | 8,3 | Te | 98 | Pb208 | 7,0 |
| Be | -0,0002622 | 0,0603 | 0,213 | 81 | Pb208 | 0,551 | 0,612 | 0,716 | 7,0 | Be | 81 | Ti | 6,9 |
| Bi | 0,00984 | 0,0197 | 0,0277 | 23 | Pd | -0,0048 | -0,0007 | 0,0062 | -317 | Ta | 78 | Th | 6,5 |
| Ca | 20140 | 21268 | 22813 | 3,4 | Pt | -0,0024 | -0,0001 | 0,0037 | -668 | Se | 46 | Rb | 6,4 |
| Cd | 0,0204 | 0,0389 | 0,0538 | 23 | Rb | 2,39 | 2,73 | 3,03 | 6,4 | S | 32 | Co | 6,4 |
| Ce | 4,84 | 5,2 | 5,96 | 3 | Re | -0,0015154 | 0,000319 | 0,00386 | 148 | Ag | 29 | Zn | 6,2 |
| Co | 1,91 | 2,08 | 2,4 | 6,4 | S | -197,2 | 1018 | 1435 | 32 | V | 28 | U | 6,2 |
| Cr | 12,1 | 13,8 | 16,9 | 4,2 | Sb | 0,0122 | 0,0329 | 0,0482 | 21 | Au | 25 | Cu | 5,6 |
| Cs | 0,114 | 0,129 | 0,148 | 5,2 | Sc | 0,637 | 0,92 | 1,17 | 13 | Li | 24 | P | 5,4 |
| Cu | 8,24 | 9,36 | 10,2 | 5,6 | Se | -0,0415573 | 0,323 | 0,571 | 46 | Bi | 23 | Ni | 5,2 |
| Fe | 2706 | 2886 | 3215 | 4,2 | Sn | 0,0513 | 0,092 | 0,145 | 17 | Cd | 23 | Cs | 5,2 |
| Ga | 0,557 | 0,678 | 0,804 | 7,1 | Sr | 115 | 124 | 129 | 3,6 | Sb | 21 | Al | 5,1 |
| Ge | -0,0658984 | 0,0268 | 0,081 | 126 | Ta | -0,0020997 | 0,00119 | 0,00348 | 78 | W | 18 | K | 4,3 |
| Hf | 0,0243 | 0,0494 | 0,0658 | 17 | Te | -0,0127631 | 0,0187 | 0,073 | 98 | Sn | 17 | Cr | 4,2 |
| Hg | 0,0389 | 0,054 | 0,0718 | 8,2 | Th | 0,659 | 0,727 | 0,821 | 6,5 | Hf | 17 | Fe | 4,2 |
| In | -0,0040796 | 0,00229 | 0,00962 | 99 | Ti | 27,3 | 30,9 | 34,5 | 6,9 | As | 17 | Ba | 3,9 |
| K | 1662 | 1779 | 1912 | 4,3 | Tl | 0,0111 | 0,0162 | 0,0222 | 11 | Sc | 13 | Y | 3,8 |
| La | 2,3 | 2,55 | 2,65 | 3,3 | U | 0,161 | 0,185 | 0,214 | 6,2 | Nb | 13 | Sr | 3,6 |
| Li | 0,423 | 0,576 | 0,753 | 24 | V | 0,405 | 3,41 | 4,58 | 28 | Mo | 12 | Ca | 3,4 |
| Mg | 1256 | 1306 | 1445 | 3,3 | W | 0,0423 | 0,0699 | 0,141 | 18 | Tl | 11 | La | 3,3 |
| Mn | 395 | 429 | 466 | 3,3 | Y | 1,48 | 1,56 | 1,68 | 3,8 | B | 9,0 | Mg | 3,3 |
| Mo | 0,174 | 0,215 | 0,259 | 12 | Zn | 22,8 | 26,2 | 29,1 | 6,2 | Pb204 | 8,6 | Mn | 3,3 |
| Na | 52,3 | 60 | 68,3 | 7,2 | Zr | 1,1 | 1,26 | 1,45 | 7,6 | Pb207 | 8,3 | Ce | 3,0 |
| Nb | 0,0314 | 0,0586 | 0,0702 | 13 | | | | | | Pb | 8,2 | | |

Table 4: Minimum, median (Q50), maximum and precision values for the Acme standard V16. Concentrations in mg/kg

| Acme V16 standard (N=28) alphabetical | | | | | | Sorted by precision | | | | | | | |
|---------------------------------------|------------|------------|---------|------------|-------------------|---------------------|----------|---------|-----------|---------|-------------|-------------------|-----------|
| Element | Min | Q50 | Max | Precision | Element | Min | Q50 | Max | Precision | Element | Precision | Element | Precision |
| | | | | | | | | | | | | | |
| Ag | 0,034 | 0,0409 | 0,048 | 9,5 | Ni | 6,5 | 8,53 | 10,3 | 11 | Be | 234648346,2 | Co | 13 |
| Al | 495 | 535 | 612 | 3,2 | P | 472 | 524 | 578 | 4,4 | Re | 1585317 | B | 12 |
| As | 1,14 | 1,65 | 2,27 | 13 | ²⁰⁴ Pb | 3,22 | 3,44 | 4,16 | 2,6 | Pd | 914 | Cr | 11 |
| Au | 0,000392 | 0,00121 | 0,00788 | 47 | ²⁰⁶ Pb | 0,0417 | 0,0461 | 0,0527 | 3,8 | In | -464 | Ni | 11 |
| B | 3,78 | 4,99 | 6,38 | 12 | ²⁰⁷ Pb | 0,81 | 0,868 | 1,05 | 3,3 | Pt | 257 | Sb | 11 |
| Ba | 1,51 | 2,18 | 2,59 | 7,0 | ²⁰⁸ Pb | 0,684 | 0,727 | 0,909 | 2,5 | S | 173 | Ce | 11 |
| Be | -0,0379628 | 0 | 0,0465 | -234648346 | Rb | 1,64 | 1,8 | 2,16 | 3,4 | Se | 161 | Fe | 11 |
| Bi | 0,00152 | 0,0105 | 0,015 | 15 | Pd | -0,0028 | 0,00023 | 0,00438 | 914 | Te | 144 | La | 10 |
| Ca | 3271 | 3468 | 3690 | 4,3 | Pt | -0,0019 | 0,000148 | 0,00112 | 257 | V | 115 | Cu | 10 |
| Cd | 0,0644 | 0,0898 | 0,153 | 18 | Re | -0,0015 | 0 | 0,00691 | 2E+06 | Sc | 88 | Cs | 8,4 |
| Ce | 0,0847 | 0,113 | 0,162 | 11 | S | -1223 | 222 | 604 | 173 | Th | 77 | Na | 8,2 |
| Co | 0,939 | 1,2 | 1,93 | 13 | ²⁰⁸ Pb | 0,0604 | 0,0765 | 0,105 | 11 | Ta | 72 | Ba | 7,0 |
| Cr | 284 | 364 | 455 | 11 | Sc | -0,0369002 | 0,207 | 0,555 | 88 | Ge | 72 | Ti | 6,6 |
| Cs | 0,0346 | 0,0394 | 0,0492 | 8,4 | Se | -0,1173872 | 0,124 | 0,458 | 161 | Hf | 66 | Zn | 6,0 |
| Cu | 6,45 | 7,5 | 20,8 | 9,8 | Sn | 0,154 | 0,239 | 1,03 | 17 | Au | 47 | Rb | 6,0 |
| Fe | 3751 | 4702 | 5669 | 11 | Sr | 10,9 | 11,5 | 12,2 | 4,6 | Ga | 41 | Sr | 4,6 |
| Ga | 0,136 | 0,256 | 0,354 | 41 | Ta | -0,000995 | 0,00104 | 0,00257 | 72 | Li | 32 | Mn | 4,6 |
| Ge | -0,0474746 | 0,0493 | 0,131 | 72 | Te | -0,0593523 | 0,0108 | 0,113 | 144 | W | 29 | P | 4,4 |
| Hf | 0 | 0,00806 | 0,0205 | 66 | Th | -0,0506307 | 0,00524 | 0,0106 | 77 | Hg | 20 | K | 4,4 |
| Hg | 0,0414 | 0,0551 | 0,0681 | 20 | Ti | 11,2 | 12,8 | 14,8 | 6,6 | Tl | 20 | Ca | 4,3 |
| In | -0,0050964 | -0,0005054 | 0,00499 | -464 | Tl | 0,00841 | 0,0126 | 0,0183 | 20 | Zr | 19 | ²⁰⁴ Pb | 3,8 |
| K | 2105 | 2313 | 2488 | 4,4 | U | -0,0008899 | 0,00288 | 0,00574 | 101 | Nb | 19 | ²⁰⁸ Pb | 3,4 |
| La | 0,0359 | 0,0504 | 0,0847 | 10 | W | -1,5032125 | 0,462 | 1,46 | 115 | Mo | 18 | ²⁰⁶ Pb | 3,3 |
| Li | 0,0281 | 0,066 | 0,13 | 32 | Y | 0,0154 | 0,0372 | 0,0702 | 29 | Cd | 18 | Mg | 3,3 |
| Mg | 544 | 584 | 663 | 3,3 | Zn | 37,8 | 44,9 | 48,6 | 6,0 | Sn | 17 | Al | 3,2 |
| Mn | 704 | 780 | 858 | 4,6 | Zr | 0,122 | 0,17 | 0,507 | 13 | Bi | 15 | Pb | 2,6 |
| Mo | 1,31 | 1,79 | 2,58 | 18 | | | | | 19 | As | 13 | ²⁰⁷ Pb | 2,5 |
| Na | 15,8 | 18,5 | 23,4 | 8,2 | | | | | | Y | 13 | | |
| Nb | 0,0768 | 0,108 | 0,149 | 19 | | | | | | | | | |

Table 5: Precision on analytical and field duplicates and also replicate analyses..

| Field duplicates (38 pairs) | | | | Analytical duplicates (38 pairs) | | | | Replicate analyses (28 reps) | | | |
|-----------------------------|-----------|---------|-----------|----------------------------------|-----------|---------|-----------|------------------------------|-----------|---------|-----------|
| Alphabetical | | Sorted | | Alphabetical | | Sorted | | Alphabetical | | Sorted | |
| Element | Precision | Element | Precision | Element | Precision | Element | Precision | Element | Precision | Element | Precision |
| Ag | 29 | Pd | -1588 | Ag | 8,1 | Pt | -39833 | Ag | 2,6 | Pd | 2187 |
| Al | 48 | Pt | -1153 | Al | 5,5 | Pd | -2229 | Al | 6,5 | Re | 324 |
| As | 34 | Au | 351 | As | 23 | Au | 305 | As | 10 | Pt | 243 |
| Au | 351 | Re | 232 | Au | 305 | Re | 180 | Au | 75 | Te | 148 |
| B | 24 | Te | 137 | B | 20 | Te | 91 | B | 22 | Au | 75 |
| Ba | 28 | V | 90 | Ba | 4,5 | Be | 71 | Ba | 2,8 | Be | 65 |
| Be | 71 | Ti | 83 | Be | 71 | In | 64 | Be | 65 | In | 53 |
| Bi | 21 | Ga | 81 | Bi | 7,9 | Hf | 31 | Bi | 7,5 | W | 32 |
| Ca | 20 | Mn | 75 | Ca | 7,3 | Ta | 31 | Ca | 2,8 | Ta | 22 |
| Cd | 21 | U | 71 | Cd | 9,1 | La | 29 | Cd | 6,4 | B | 22 |
| Ce | 66 | Be | 71 | Ce | 12 | Ge | 27 | Ce | 6,1 | Ge | 21 |
| Co | 48 | In | 67 | Co | 18 | As | 23 | Co | 12 | Hf | 19 |
| Cr | 60 | Ce | 66 | Cr | 5,2 | W | 22 | Cr | 6,9 | Mn | 17 |
| Cs | 57 | Y | 63 | Cs | 5,2 | U | 21 | Cs | 3,9 | Se | 17 |
| Cu | 44 | Li | 61 | Cu | 8,1 | B | 20 | Cu | 11 | Mg | 12 |
| Fe | 51 | Cr | 60 | Fe | 5,2 | Y | 19 | Fe | 8,4 | Co | 12 |
| Ga | 81 | La | 59 | Ga | 7,3 | Co | 18 | Ga | 7,0 | Cu | 11 |
| Ge | 36 | Nb | 59 | Ge | 27 | Se | 14 | Ge | 21 | As | 10 |
| Hf | 40 | Cs | 57 | Hf | 31 | Th | 14 | Hf | 19 | Sb | 10 |
| Hg | 21 | Fe | 51 | Hg | 8,0 | Mn | 14 | Hg | 5,4 | Mo | 9,2 |
| In | 67 | Th | 50 | In | 64 | Ce | 12 | In | 53 | V | 8,7 |
| K | 14 | Zr | 49 | K | 3,8 | Mo | 12 | K | 2,5 | Fe | 8,4 |
| La | 59 | Co | 48 | La | 29 | Ni | 11 | La | 6,4 | S | 8,2 |
| Li | 61 | Al | 48 | Li | 10 | Tl | 11 | Li | 7,9 | Nb | 8,1 |
| Mg | 31 | Cu | 44 | Mg | 3,9 | Li | 10 | Mg | 12 | Zr | 8,0 |
| Mn | 75 | Ta | 42 | Mn | 14 | Sc | 10 | Mn | 17 | Li | 7,9 |
| Mo | 27 | Hf | 40 | Mo | 12 | Cd | 9,1 | Mo | 9,2 | Tl | 7,6 |
| Na | 8,3 | Ge | 36 | Na | 5,6 | Ag | 8,1 | Na | 5,4 | Bi | 7,5 |
| Nb | 59 | As | 34 | Nb | 6,1 | Cu | 8,1 | Nb | 8,1 | Sc | 7,5 |
| Ni | 27 | Se | 31 | Ni | 11 | Hg | 8,0 | Ni | 5,5 | Th | 7,5 |
| P | 13 | Mg | 31 | P | 4,6 | Bi | 7,9 | P | 2,7 | U | 7,2 |
| Pb | 19 | Ag | 29 | Pb | 4,5 | Ca | 7,3 | Pb | 2,5 | Ga | 7,0 |
| Pd | -1588 | Tl | 29 | Pd | -2229 | Ga | 7,3 | Pd | 2187 | Cr | 6,9 |
| Pt | -1153 | Sc | 28 | Pt | -39833 | Sn | 7,1 | Pt | 243 | Al | 6,5 |
| Rb | 26 | Ba | 28 | Rb | 5,5 | Zr | 6,8 | Rb | 5,0 | Cd | 6,4 |
| Re | 232 | Ni | 27 | Re | 180 | V | 6,7 | Re | 324 | La | 6,4 |
| S | 22 | Mo | 27 | S | 6,2 | Ti | 6,4 | S | 8,2 | Ce | 6,1 |
| Sb | 21 | W | 27 | Sb | 5,7 | S | 6,2 | Sb | 10 | Ni | 5,5 |
| Sc | 28 | Rb | 26 | Sc | 10 | Nb | 6,1 | Sc | 7,5 | Hg | 5,4 |
| Se | 31 | Zn | 25 | Se | 14 | Sb | 5,7 | Se | 17 | Na | 5,4 |
| Sn | 23 | B | 24 | Sn | 7,1 | Na | 5,6 | Sn | 5,2 | Sn | 5,2 |
| Sr | 20 | Sn | 23 | Sr | 5,2 | Al | 5,5 | Sr | 2,0 | Rb | 5,0 |
| Ta | 42 | S | 22 | Ta | 31 | Rb | 5,5 | Ta | 22 | Ti | 5,0 |
| Te | 137 | Cd | 21 | Te | 91 | Cr | 5,2 | Te | 148 | Zn | 4,5 |
| Th | 50 | Bi | 21 | Th | 14 | Cs | 5,2 | Th | 7,5 | Cs | 3,9 |
| Ti | 83 | Hg | 21 | Ti | 6,4 | Fe | 5,2 | Ti | 5,0 | Y | 3,5 |
| Tl | 29 | Sb | 21 | Tl | 11 | Sr | 5,2 | Tl | 7,6 | Ba | 2,8 |
| U | 71 | Ca | 20 | U | 21 | Zn | 5,0 | U | 7,2 | Ca | 2,8 |
| V | 90 | Sr | 20 | V | 6,7 | P | 4,6 | V | 8,7 | P | 2,7 |
| W | 27 | Pb | 19 | W | 22 | Ba | 4,5 | W | 32 | Ag | 2,6 |
| Y | 63 | K | 14 | Y | 19 | Pb | 4,5 | Y | 3,5 | K | 2,5 |
| Zn | 25 | P | 13 | Zn | 5,0 | Mg | 3,9 | Zn | 4,5 | Pb | 2,5 |
| Zr | 49 | Na | 8,3 | Zr | 6,8 | K | 3,8 | Zr | 8,0 | Sr | 2,0 |

Table 6: Statistical results (minimum, mean, maximum and standard deviation) for the NIST standard

| MINS 2013 organic soil analyses | | | | | | Reference values* | |
|-----------------------------------|---------|----------|----------|----------|----------|-------------------|----------------|
| Pb ratio | | Min. | Mean | Max. | 2*StDev | Mean value | ± |
| $^{204}\text{Pb}/^{206}\text{Pb}$ | SRM 981 | 0.055346 | 0.058771 | 0.060557 | 0.002797 | 0.059042 | $\pm 0,000037$ |
| $^{207}\text{Pb}/^{206}\text{Pb}$ | SRM 981 | 0.891885 | 0.916921 | 0.964005 | 0.032714 | 0.914640 | $\pm 0,00033$ |
| $^{208}\text{Pb}/^{206}\text{Pb}$ | SRM 981 | 1.971519 | 2.158194 | 2.282513 | 0.138801 | 2.168100 | $\pm 0,0008$ |
| $^{204}\text{Pb}/^{206}\text{Pb}$ | SRM 983 | 0.000318 | 0.000375 | 0.000443 | 0.000058 | 0.000371 | $\pm 0,00002$ |
| $^{207}\text{Pb}/^{206}\text{Pb}$ | SRM 983 | 0.066639 | 0.071622 | 0.073890 | 0.001662 | 0.071201 | $\pm 0,00004$ |
| $^{208}\text{Pb}/^{206}\text{Pb}$ | SRM 983 | 0.011815 | 0.014490 | 0.015871 | 0.003116 | 0.013619 | $\pm 0,000024$ |

*Overall limits of error are based on 95 % confidence limits for the mean of the ratio measurements and on allowances for the known sources of possible systematic error.

Table 7: Statistical results (minimum, 25 percentile, median, mean, 75-percentile, maximum, standard deviation and precision) for previous and present analyses of the Reimann et. al. (2007) study ($N=10$).

| Pb ratio | | Min | 25 % | Median | Mean | 75 % | Max | 2xSD | Prec (%) |
|-----------------------------------|----------|--------|--------|--------|--------|--------|--------|--------|----------|
| $^{206}\text{Pb}/^{207}\text{Pb}$ | previous | 1.1406 | 1.1519 | 1.1526 | 1.1642 | 1.1593 | 1.2300 | 0.0546 | 0.7 |
| | present | 1.1379 | 1.1503 | 1.1593 | 1.1659 | 1.1743 | 1.2150 | 0.0515 | |
| $^{206}\text{Pb}/^{208}\text{Pb}$ | previous | 0.4734 | 0.4740 | 0.4743 | 0.4775 | 0.4754 | 0.4952 | 0.0145 | 1.8 |
| | present | 0.4595 | 0.4792 | 0.4850 | 0.4809 | 0.4859 | 0.4881 | 0.0175 | |
| $^{208}\text{Pb}/^{207}\text{Pb}$ | previous | 2.4090 | 2.4290 | 2.4304 | 2.4373 | 2.4386 | 2.4836 | 0.0412 | 1.5 |
| | present | 2.3412 | 2.3803 | 2.4002 | 2.4249 | 2.4669 | 2.5431 | 0.1337 | |

Table 8: Statistical results (minimum, 25 percentile, median, mean, 75-percentile, maximum standard deviation and precision) for previous and present analyses of the Jensen & Finne (2006) study ($N=5$).

| Pb ratio | | Min | 25 % | Median | Mean | 75 % | Max | 2xSD | Prec (%) |
|-----------------------------------|----------|--------|--------|--------|--------|--------|--------|--------|----------|
| $^{206}\text{Pb}/^{207}\text{Pb}$ | previous | 1.1486 | 1.1510 | 1.1578 | 1.1603 | 1.1580 | 1.1861 | 0.0300 | 1.2 |
| | present | 1.1266 | 1.1347 | 1.1429 | 1.1452 | 1.1580 | 1.1638 | 0.0312 | |
| $^{206}\text{Pb}/^{208}\text{Pb}$ | previous | 0.4747 | 0.4755 | 0.4765 | 0.4779 | 0.4771 | 0.4858 | 0.0090 | 1.5 |
| | present | 0.4596 | 0.4643 | 0.4706 | 0.4708 | 0.4790 | 0.4807 | 0.0182 | |
| $^{208}\text{Pb}/^{207}\text{Pb}$ | previous | 2.4168 | 2.4211 | 2.4285 | 2.4288 | 2.4362 | 2.4415 | 0.0205 | 0.5 |
| | present | 2.4091 | 2.4288 | 2.4294 | 2.4324 | 2.4436 | 2.4511 | 0.0322 | |

Table 9: Statistical parameters for the mapped data. Humus <2mm, aqua regia extraction on 5g sample material. N=752, concentrations in mg/kg

| Ele. | n<PDL | MDL | PDL | Min | Q2 | Q5 | Q10 | Q25 | Q50 | Q75 | Q90 | Q95 | Q98 | Max | Power |
|------|-------|--------|--------|---------|---------|---------|---------|---------|---------|--------|--------|--------|--------|--------|-------|
| Ag | 0 | 0.002 | 0.002 | 0.029 | 0.047 | 0.057 | 0.068 | 0.11 | 0.20 | 0.44 | 0.86 | 1.14 | 1.76 | 5.87 | 2.3 |
| Al | 0 | 100 | 100 | 304 | 465 | 572 | 734 | 1087 | 2030 | 4221 | 7484 | 10435 | 14035 | 43010 | 2.2 |
| As | 110 | 0.1 | 0.3 | <0.3 | <0.3 | <0.3 | <0.3 | 0.5 | 0.8 | 1.3 | 1.9 | 2.7 | 5.7 | 112 | 2.9 |
| Au | 445 | 0.0002 | 0.0005 | <0.0005 | <0.0005 | <0.0005 | <0.0005 | <0.0005 | <0.0005 | 0.0008 | 0.0013 | 0.002 | 0.002 | 0.02 | 2 |
| B | 12 | 1 | 0.5 | <0.50 | 0.6 | 0.7 | 1.0 | 1.3 | 1.8 | 2.4 | 3.0 | 3.7 | 4.3 | 6.5 | 1.4 |
| Ba | 0 | 0.1 | 0.1 | 6.54 | 11.0 | 13.1 | 16.5 | 23.9 | 35.5 | 50.1 | 70.4 | 89.6 | 114 | 286 | 1.6 |
| Be | 747 | 0.1 | 1 | <1.0 | <1.0 | <1.0 | <1.0 | <1.0 | <1.0 | <1.0 | <1.0 | <1.0 | <1.0 | <1.0 | 4.7 |
| Bi | 0 | 0.02 | 0.02 | 0.02 | 0.04 | 0.06 | 0.07 | 0.10 | 0.13 | 0.17 | 0.21 | 0.23 | 0.28 | 0.47 | 1.3 |
| Ca | 0 | 100 | 100 | 403 | 830 | 1148 | 1438 | 2086 | 2681 | 3446 | 4425 | 5344 | 8115 | 21802 | 1.7 |
| Cd | 0 | 0.01 | 0.05 | 0.06 | 0.12 | 0.19 | 0.25 | 0.36 | 0.51 | 0.71 | 0.92 | 1.1 | 1.3 | 1.9 | 1.5 |
| Ce | 0 | 0.01 | 0.01 | 0.48 | 0.72 | 0.93 | 1.3 | 2.2 | 5.1 | 12 | 25 | 38 | 68 | 404 | 2.9 |
| Co | 0 | 0.01 | 0.01 | 0.21 | 0.33 | 0.44 | 0.57 | 0.85 | 1.5 | 2.7 | 5.1 | 8.1 | 19 | 93 | 2.7 |
| Cr | 0 | 0.1 | 0.1 | 1.1 | 1.3 | 1.5 | 1.7 | 2.1 | 3.0 | 6.3 | 13 | 21 | 34 | 1112 | 3 |
| Cs | 0 | 0.005 | 0.005 | 0.026 | 0.053 | 0.068 | 0.090 | 0.139 | 0.266 | 0.508 | 0.974 | 1.30 | 1.81 | 4.31 | 2.2 |
| Cu | 0 | 0.01 | 0.01 | 2.7 | 4.2 | 4.8 | 5.5 | 6.4 | 7.9 | 10 | 14 | 19 | 34 | 305 | 2 |
| Fe | 0 | 10 | 10 | 376 | 601 | 704 | 930 | 1490 | 3003 | 6168 | 13293 | 18635 | 28188 | 183346 | 2.7 |
| Ga | 0 | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.3 | 0.5 | 0.8 | 1.6 | 3.0 | 4.2 | 6.1 | 11 | 1.9 |
| Ge | 54 | 0.01 | 0.02 | <0.02 | <0.02 | <0.02 | 0.03 | 0.07 | 0.14 | 0.22 | 0.31 | 0.38 | 0.44 | 0.64 | 1.8 |
| Hf | 40 | 0.001 | 0.006 | <0.006 | 0.003 | 0.003 | 0.009 | 0.014 | 0.023 | 0.041 | 0.066 | 0.087 | 0.12 | 0.21 | 1.8 |
| Hg | 0 | 0.001 | 0.001 | 0.059 | 0.093 | 0.12 | 0.14 | 0.18 | 0.21 | 0.26 | 0.32 | 0.36 | 0.40 | 1.0 | 1.2 |
| In | 671 | 0.02 | 0.02 | <0.02 | <0.02 | <0.02 | <0.02 | <0.02 | <0.02 | <0.02 | 0.02 | 0.03 | 0.03 | 0.05 | 0.7 |
| K | 0 | 100 | 100 | 282 | 477 | 536 | 598 | 711 | 851 | 1009 | 1226 | 1428 | 1674 | 2527 | 1 |
| La | 0 | 0.01 | 0.01 | 0.21 | 0.31 | 0.42 | 0.57 | 1.00 | 2.4 | 5.3 | 12 | 20 | 39 | 174 | 2.9 |
| Li | 138 | 0.01 | 0.1 | <0.1 | <0.1 | <0.1 | <0.1 | 0.1 | 0.2 | 0.6 | 2.6 | 5.0 | 9.4 | 47 | 3 |
| Mg | 0 | 10 | 10 | 302 | 591 | 794 | 912 | 1112 | 1401 | 1761 | 2449 | 3669 | 5257 | 35752 | 2.1 |
| Mn | 0 | 1 | 1 | 7.5 | 11 | 14 | 18 | 30 | 58 | 117 | 342 | 636 | 1270 | 5254 | 2.8 |
| Mo | 0 | 0.01 | 0.01 | 0.09 | 0.16 | 0.19 | 0.22 | 0.28 | 0.39 | 0.62 | 0.98 | 1.3 | 2.0 | 12 | 2.1 |
| Na | 0 | 10 | 10 | 28 | 52 | 64 | 75 | 103 | 153 | 257 | 359 | 417 | 484 | 789 | 1.5 |
| Nb | 0 | 0.01 | 0.01 | 0.03 | 0.05 | 0.07 | 0.10 | 0.18 | 0.42 | 1.0 | 2.0 | 2.8 | 3.5 | 6.8 | 2.3 |
| Ni | 0 | 0.1 | 0.1 | 0.9 | 1.2 | 1.4 | 1.6 | 2.2 | 3.2 | 5.3 | 9.4 | 14 | 22 | 361 | 2.6 |
| P | 0 | 10 | 10 | 200 | 386 | 462 | 534 | 644 | 775 | 951 | 1152 | 1276 | 1447 | 2141 | 1 |
| Pb | 0 | 0.01 | 0.01 | 3.1 | 9.7 | 12.7 | 15.8 | 20.2 | 27 | 34.7 | 43.8 | 50.9 | 64.4 | 3511 | 3.1 |
| Pd | 748 | 0.002 | 0.01 | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | 0.018 | 0.6 |
| Pt | 739 | 0.001 | 0.002 | <0.002 | <0.002 | <0.002 | <0.002 | <0.002 | <0.002 | <0.002 | <0.002 | <0.002 | <0.002 | 0.024 | 1.4 |
| Rb | 0 | 0.1 | 0.1 | 0.84 | 1.4 | 1.7 | 2.1 | 2.8 | 4.2 | 6.2 | 9.5 | 13 | 17 | 30 | 1.6 |
| Re | 605 | 0.001 | 0.001 | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | <0.001 | 0.002 | 0.002 | 0.003 | 0.011 | 1.3 |
| S | 13 | 100 | 50 | <50 | 64 | 394 | 662 | 1048 | 1441 | 1854 | 2262 | 2464 | 2888 | 4008 | 2.2 |
| Sb | 0 | 0.02 | 0.02 | 0.05 | 0.13 | 0.16 | 0.19 | 0.25 | 0.33 | 0.43 | 0.53 | 0.60 | 0.68 | 123 | 3.4 |
| Sc | 9 | 0.1 | 0.15 | <0.15 | 0.21 | 0.30 | 0.40 | 0.57 | 0.81 | 1.2 | 1.7 | 2.2 | 3.5 | 11 | 2.2 |
| Se | 16 | 0.1 | 0.15 | <0.15 | <0.15 | 0.23 | 0.32 | 0.55 | 0.90 | 1.3 | 1.8 | 2.3 | 2.9 | 5.3 | 1.8 |
| Sn | 1 | 0.02 | 0.15 | <0.15 | 0.30 | 0.37 | 0.43 | 0.58 | 0.75 | 0.95 | 1.1 | 1.3 | 1.5 | 10 | 2.1 |
| Sr | 0 | 0.5 | 0.5 | 4.44 | 10.3 | 13.9 | 16.9 | 22.4 | 29.9 | 37.6 | 45.9 | 52.2 | 59.1 | 96.9 | 1.3 |
| Ta | 16 | 0.001 | 0.001 | <0.001 | <0.001 | 0.002 | 0.003 | 0.006 | 0.011 | 0.017 | 0.026 | 0.032 | 0.042 | 0.080 | 2.2 |
| Te | 753 | 0.02 | 0.2 | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 | <0.2 | 0.2 | 0.3 |
| Th | 4 | 0.01 | 0.005 | <0.005 | 0.030 | 0.048 | 0.066 | 0.11 | 0.20 | 0.38 | 0.78 | 1.2 | 1.9 | 7.6 | 3.5 |
| Ti | 0 | 1 | 1 | 21 | 30 | 40 | 53 | 87 | 161 | 316 | 622 | 914 | 1437 | 4437 | 2.3 |
| TI | 0 | 0.02 | 0.01 | 0.02 | 0.03 | 0.04 | 0.05 | 0.06 | 0.08 | 0.11 | 0.14 | 0.16 | 0.20 | 0.55 | 1.5 |
| U | 0 | 0.01 | 0.01 | 0.02 | 0.03 | 0.04 | 0.05 | 0.09 | 0.20 | 0.49 | 1.3 | 2.2 | 4.7 | 24 | 3.2 |
| V | 28 | 2 | 0.2 | <0.2 | <0.2 | 0.3 | 0.7 | 1.5 | 2.9 | 6.6 | 14 | 20 | 32 | 80 | 2.9 |
| W | 0 | 0.1 | 0.01 | 0.01 | 0.03 | 0.04 | 0.04 | 0.06 | 0.08 | 0.11 | 0.16 | 0.20 | 0.30 | 3.0 | 2.3 |
| Y | 0 | 0.001 | 0.001 | 0.10 | 0.16 | 0.21 | 0.27 | 0.47 | 1.1 | 2.5 | 5.1 | 8.1 | 14 | 65 | 2.8 |
| Zn | 0 | 0.1 | 0.1 | 3.38 | 12.8 | 17.0 | 21.4 | 28.8 | 38.3 | 48.8 | 61.8 | 71.7 | 87.1 | 247 | 1.9 |
| Zr | 0 | 0.01 | 0.01 | 0.07 | 0.23 | 0.27 | 0.33 | 0.50 | 0.77 | 1.2 | 2.2 | 3.0 | 3.9 | 11 | 2.2 |


| Pb ratio | Min | Q2 | Q5 | Q10 | Q25 | Q50 | Q75 | Q90 | Q95 | Q98 | Max | Powers |
|--------------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| ²⁰⁸ Pb/ ²⁰⁶ Pb | 1.8595 | 1.9974 | 2.0158 | 2.0333 | 2.0676 | 2.1020 | 2.1368 | 2.1677 | 2.1912 | 2.2115 | 2.2418 | 0.1 |
| ²⁰⁶ Pb/ ²⁰⁷ Pb | 1.0399 | 1.1136 | 1.1190 | 1.1266 | 1.1379 | 1.1512 | 1.1650 | 1.1781 | 1.1894 | 1.2054 | 1.2584 | 0.1 |


Table 10: Comparison between Nord-Trøndelag & Fosen survey with Njåstad et. al. (1994), where the greatest value between the datasets is underlined. Statistics from other organic soil surveys covering parts of the MINS area from Ryghaug (1980) and Sæther (1985) (see Figure 1) are also given.


| | MINS 2013 (N=754) | | | Njåstad et. al, 1994 (N=527) | | | Sæther, 1985 (N=217) | | | Ryghaug, 1980 (N=1244) | | |
|------|----------------------|--------------|---------------|---------------------------------|-------|-------|-------------------------|-------|-------|---------------------------|------|-------|
| Ele. | Q50 | Q98 | Max | Q50 | Q98 | Max | Q50 | Q98 | Max | Q50 | Q98 | Max |
| Ag | 0.20 | 1.76 | <u>5.87</u> | 0.3 | 1 | 2.2 | 0.3 | 1.1 | 2.5 | 0.3 | 1.3 | 5.4 |
| Al | 2030 | 14035 | <u>43010</u> | 1700 | 15400 | 27700 | 1832 | 8854 | 18336 | | | |
| As | 0.8 | 5.7 | 112 | | | | | | | | | |
| Au | <0.0005 | 0.002 | 0.02 | | | | | | | | | |
| B | 1.8 | 4.3 | 6.5 | 2.6 | 7 | 58.7 | 5 | 14.7 | 26.2 | | | |
| Ba | 35.5 | 114 | 286 | 45.8 | 238 | 483 | 31.4 | 109 | 162 | | | |
| Be | <1.0 | <1.0 | <u>4.7</u> | 0.2 | 2 | 3 | 0 | 0.2 | 0.4 | | | |
| Bi | 0.13 | 0.28 | 0.47 | | | | | | | | | |
| Ca | 2681 | 8115 | <u>21802</u> | 2200 | 8050 | 18600 | 3003 | 12595 | 21433 | | | |
| Cd | 0.51 | 1.3 | 1.9 | 0.8 | 3 | 12.4 | 0.4 | 1.27 | 1.6 | 0.5 | 1.7 | 14.6 |
| Ce | <u>5.1</u> | <u>68</u> | <u>404</u> | 4.5 | 51 | 130 | | | | | | |
| Co | 1.5 | 19 | <u>93</u> | 1.9 | 13 | 29.8 | 1.1 | 9.61 | 21.1 | 4 | 40 | 131 |
| Cr | 3.0 | 34 | <u>1112</u> | | | | 2.2 | 22.3 | 49.8 | 5 | 68.1 | 268 |
| Cs | 0.266 | 1.81 | 4.31 | | | | | | | | | |
| Cu | 7.9 | 34 | 305 | | | | 5.7 | 18.7 | 24.8 | 9 | 55.4 | 2047 |
| Fe | <u>3003</u> | <u>28188</u> | <u>183346</u> | 2100 | 19600 | 49200 | 1738 | 13856 | 30369 | 0.27 | 3.49 | 16.6 |
| Ga | 0.8 | 6.1 | 11 | | | | | | | | | |
| Ge | 0.14 | 0.44 | 0.64 | | | | | | | | | |
| Hf | 0.023 | 0.12 | 0.21 | | | | | | | | | |
| Hg | 0.21 | 0.40 | 1.0 | 3 | 32 | 189 | | | | | | |
| In | <0.02 | 0.03 | 0.05 | 7 | 58 | 1300 | | | | | | |
| K | 851 | 1674 | 2527 | 820 | 1950 | 3400 | 1119 | 2563 | 6017 | | | |
| La | <u>2.4</u> | <u>39</u> | 174 | 2 | 33 | 198 | | | | | | |
| Li | 0.2 | 9.4 | <u>47</u> | 0.5 | 10 | 28.7 | 0.5 | 5.87 | 15 | | | |
| Mg | <u>1401</u> | <u>5257</u> | <u>35752</u> | 1100 | 4550 | 10900 | 1360 | 4982 | 11575 | | | |
| Mn | 58 | 1270 | <u>5254</u> | 60 | 1100 | 10000 | 139 | 1175 | 3602 | 145 | 2475 | 24411 |
| Mo | 0.39 | 2.0 | <u>12</u> | 0.6 | 3 | 9.5 | 0.4 | 1.44 | 2.6 | 0 | 2 | 9 |
| Na | 153 | 484 | 789 | 130 | 485 | 1700 | 304 | 821 | 1504 | | | |
| Nb | 0.42 | 3.5 | 6.8 | | | | | | | | | |
| Ni | 3.2 | 22 | 361 | 3.2 | 40 | 487 | 2.1 | 11.6 | 31.4 | 5 | 48.1 | 208 |
| P | <u>775</u> | <u>1447</u> | 2141 | 760 | 1400 | 2500 | 671 | 1413 | 2529 | | | |
| Pb | <u>27</u> | 64.4 | <u>3511</u> | 31.7 | 201 | 488 | 14.3 | 31.4 | 106 | 15 | 48 | 521 |
| Pd | <0.01 | <0.01 | 0.018 | | | | | | | | | |
| Pt | <0.002 | <0.002 | 0.024 | | | | | | | | | |
| Rb | 4.2 | 17 | 30 | | | | | | | | | |
| Re | <0.001 | 0.003 | 0.011 | | | | | | | | | |
| S | 1441 | 2888 | 4008 | | | | | | | | | |
| Sb | 0.33 | 0.68 | 123 | | | | | | | | | |
| Sc | <u>0.81</u> | <u>3.5</u> | <u>11</u> | 0.5 | 3 | 10.8 | | | | | | |
| Se | 0.90 | 2.9 | 5.3 | | | | | | | | | |
| Sn | 0.75 | 1.5 | 10 | | | | | | | | | |
| Sr | <u>29.9</u> | 59.1 | 96.9 | 23.7 | 70 | 103 | | | | | | |
| Ta | 0.011 | 0.042 | 0.080 | | | | | | | | | |
| Te | <0.2 | <0.2 | 0.2 | | | | | | | | | |
| Th | 0.20 | 1.9 | 7.6 | | | | | | | | | |
| Ti | <u>161</u> | <u>1437</u> | <u>4437</u> | 110 | 845 | 1700 | | | | | | |
| TI | 0.08 | 0.20 | 0.55 | | | | | | | | | |
| U | 0.20 | 4.7 | 24 | | | | | | | | | |
| V | 2.9 | 32 | 80 | 6.2 | 40 | 99.3 | 4.5 | 33 | 80 | 11 | 100 | 245 |
| W | 0.08 | 0.30 | 3.0 | | | | | | | | | |
| Y | 1.1 | 14 | 65 | | | | | | | | | |
| Zn | 38.3 | 87.1 | 247 | 41.8 | 156 | 1600 | 32.7 | 95.4 | 312 | 35 | 127 | 1252 |
| Zr | 0.77 | 3.9 | <u>11</u> | 0.8 | 3 | 9.4 | 0.5 | 4.04 | 6.5 | | | |


Appendix 1: Cumulative frequency diagrams for all elements


Please note that readings below the practical detection limit are set to half of the practical detection limit value























Appendix 2: Maps of geochemistry in organic soil in Nord-Trøndelag and the Fosen peninsula





<2mm Humus

Silver AR-extract

ppmAg


- 1,15 - 5,87
● 0,45 - 1,14
• 0,12 - 0,44
○ 0,07 - 0,11
○ 0,03 - 0,06






<2mm Humus
Aluminium AR-extract
ppmAl

■ 10428 - 43010
● 4213 - 10427
• 1091 - 4212
○ 574 - 1090
○ 304 - 573





<2mm Humus

Arsenic AR-extract

ppmAs


■ 2,8 - 112,4


● 1,4 - 2,7

• 0,6 - 1,3

○ 0,2 - 0,5

○ - 0,1







<2mm Humus

Gold AR-extract

ppm Au

- 0,0018 - 0,0240
● 0,0009 - 0,0017
• <0,0005 - 0,0008
○ < 0,0005
○ <0,0005







<2mm Humus

Boron AR-extract

ppmB

- 3,8 - 6,5
● 2,5 - 3,7
• 1,4 - 2,4
○ 0,8 - 1,3
○ 0,1 - 0,7







<2mm Humus

Barium AR-extract

ppmBa

- 89,5 - 286,4
● 50,2 - 89,4
• 23,9 - 50,1
○ 13,2 - 23,8
○ 6,5 - 13,1








<2mm Humus

Bismut AR-extract

ppmBi






<2mm Humus

Calcium AR-extract

ppmCa

- 5342 - 21802
● 3450 - 5341
• 2085 - 3449
○ 1151 - 2084
○ 403 - 1150




0 6 12 18 24 30 36 42 48 54 60
km


<2mm Humus

Cadmium AR-extract

ppmCd

- 1,06 - 1,91
● 0,71 - 1,05
• 0,37 - 0,70
○ 0,20 - 0,36
○ 0,05 - 0,19







<2mm Humus

Cerium AR-extract

ppmCe

- 38,0 - 403,8
● 11,7 - 37,9
• 2,3 - 11,6
○ 1,0 - 2,2
○ 0,5 - 0,9






<2mm Humus

Cobalt AR-extract

ppmCo


- 8,2 - 93,4
● 2,8 - 8,1
• 0,9 - 2,7
○ 0,5 - 0,8
○ 0,2 - 0,4




0 6 12 18 24 30 36 42 48 54 60
km

<2mm Humus
Chromium AR-extract
ppmCr
5-25-75-95-100 %ile

- 21,1 - 1112,4
- 6,4 - 21,0
- 2,2 - 6,3
- 1,6 - 2,1
- 1,1 - 1,5







<2mm Humus

Cesium AR-extract

ppmCs

- 5-25-75-95-100 %ile
- 1,31 - 4,31
 - 0,52 - 1,30
 - 0,15 - 0,51
 - 0,08 - 0,14
 - 0,03 - 0,07







<2mm Humus

Copper AR-extract

ppmCu

- 18,7 - 305,5
● 10,1 - 18,6
• 6,5 - 10,0
○ 4,9 - 6,4
○ 2,7 - 4,8







<2mm Humus

Iron AR-extract

ppmFe

- 18616 - 183346
● 6140 - 18615
• 1493 - 6139
○ 705 - 1492
○ 376 - 704






<2mm Humus

Gallium AR-extract

ppmGa

- 4,16 - 10,59
● 1,59 - 4,15
• 0,47 - 1,58
○ 0,27 - 0,46
○ 0,12 - 0,26




0 6 12 18 24 30 36 42 48 54 60
km


<2mm Humus
Germanium AR-extract

ppmGe

- 0,39 - 0,63
- 0,23 - 0,38
- 0,08 - 0,22
- 0,03 - 0,07
- < 0,02

N







<2mm Humus

Hafnium AR-extract

ppmHf

- 0,087 - 0,211
- 0,042 - 0,086
- 0,015 - 0,041
- 0,007 - 0,014
- - 0,006







<2mm Humus

Mercury AR-extract

ppmHg

- 0,37 - 1,00
● 0,27 - 0,36
• 0,18 - 0,26
○ 0,13 - 0,17
○ 0,06 - 0,12






<2mm Humus

Potassium AR-extract

ppmK


- 1429 - 2527
● 1010 - 1428
• 712 - 1009
○ 537 - 711
○ 282 - 536




0 6 12 18 24 30 36 42 48 54 60
km

<2mm Humus
Lanthanum AR-extract
ppmLa

- 20,4 - 173,7
● 5,4 - 20,3
• 1,1 - 5,3
○ 0,5 - 1,0
○ 0,2 - 0,4







<2mm Humus

Lithium AR-extract

ppmLi

- 5,1 - 47,2
- 0,7 - 5,0
- 0,2 - 0,6
- 0,2 - 0,1
- 0,0 - 0,1







<2mm Humus

Magnesium AR-extract

ppmMg

- 3645 - 35752
● 1761 - 3644
• 1107 - 1760
○ 796 - 1106
○ 302 - 795







<2mm Humus
Manganese AR-extract
ppmMn

5-25-75-95-100 %ile

- 637 - 5254
- 118 - 636
- 30 - 117
- 15 - 29
- 7 - 14







<2mm Humus

ppmMo

- 1,35 - 11,49
- 0,63 - 1,34
- 0,29 - 0,62
- 0,20 - 0,28
- 0,09 - 0,19

5-25-75-95-100 %ile








<2mm Humus

Sodium AR-extract

ppmNa







<2mm Humus

Niobium AR-extract

ppmNb

- 2,84 - 6,78
- 1,02 - 2,83
- 0,19 - 1,01
- 0,08 - 0,18
- 0,03 - 0,07






<2mm Humus

Nickel AR-extract

ppmNi


- 13,8 - 361,0
● 5,3 - 13,7
• 2,3 - 5,2
○ 1,5 - 2,2
○ 0,9 - 1,4




0 6 12 18 24 30 36 42 48 54 60
km

<2mm Humus
Phosphorus AR-extract
ppmP

- 1277 - 2141
● 952 - 1276
• 645 - 951
○ 464 - 644
○ 200 - 463







<2mm Humus

Lead AR-extract

ppmPb

- 51,0 - 3511,4
● 34,9 - 50,9
• 20,4 - 34,8
○ 12,8 - 20,3
○ 3,1 - 12,7







<2mm Humus

Rubidium AR-extract

ppmRb

- 13,0 - 29,8
● 6,3 - 12,9
• 2,9 - 6,2
○ 1,8 - 2,8
○ 0,8 - 1,7







<2mm Humus

Sulphur AR-extract

ppmS

- 2464 - 4008
● 1855 - 2463
• 1048 - 1854
○ 396 - 1047
○ - 395








<2mm Humus

Antimony AR-extract

ppmSb







<2mm Humus

Scandium AR-extract

ppmSc

- 2,16 - 10,73
● 1,19 - 2,15
• 0,58 - 1,18
○ 0,31 - 0,57
○ 0,00 - 0,30








<2mm Humus

Selenium AR-extract

ppmSe








<2mm Humus

Tin AR-extract

ppmSn







<2mm Humus
Strontium AR-extract

ppmSr

- 52,2 - 96,9
● 37,8 - 52,1
• 22,5 - 37,7
○ 14,1 - 22,4
○ 4,4 - 14,0








<2mm Humus

Tantalum AR-extract

ppmTa







<2mm Humus

Thorium AR-extract

ppmTh

- 1,24 - 7,64
- 0,38 - 1,24
- 0,11 - 0,38
- 0,05 - 0,11
- < 0,05







<2mm Humus

Titanium AR-extract

ppmTi

- 914 - 4437
● 317 - 913
• 88 - 316
○ 41 - 87
○ 21 - 40







<2mm Humus

Thallium AR-extract

ppmTl

- 0,17 - 0,55
- 0,12 - 0,16
- 0,07 - 0,11
- 0,05 - 0,06
- 0,02 - 0,04






<2mm Humus

Uranium AR-extract

ppmU


- 2,16 - 23,86
● 0,50 - 2,15
• 0,10 - 0,49
○ 0,05 - 0,09
○ 0,02 - 0,04




0 6 12 18 24 30 36 42 48 54 60
km

<2mm Humus
Vanadium AR-extract
ppmV

■ 20,3 - 80,4
● 6,6 - 20,2
• 1,6 - 6,5
○ 0,4 - 1,5
○ < 0,3








<2mm Humus

Tungsten AR-extract

ppmW







<2mm Humus

Yttrium AR-extract

ppmY

- 8,2 - 65,0
● 2,6 - 8,1
• 0,6 - 2,5
○ 0,3 - 0,5
○ 0,1 - 0,2








<2mm Humus

Zinc AR-extract

ppmZn







<2mm Humus

Zirconium AR-extract

ppmZr

- 2,97 - 10,59
● 1,25 - 2,96
• 0,51 - 1,24
○ 0,28 - 0,50
○ 0,07 - 0,27







<2mm Humus

206_207Lead AR-extract

$^{206}\text{Pb}/^{207}\text{Pb}$

- 1,1909 - 1,2584
● 1,1643 - 1,1908
• 1,1465 - 1,1642
○ 1,1278 - 1,1464
○ 1,0399 - 1,1277





<2mm Humus

208_206Lead AR-extract

$^{208}\text{Pb}/^{206}\text{Pb}$

- 2,1693 - 2,2418
● 2,1219 - 2,1692
• 2,0825 - 2,1218
○ 2,0375 - 2,0824
○ 1,8595 - 2,0374

